## Citations

- 1. Warshafsky S, Kamer RS, Sivak SL. Effect of garlic on total serum cholesterol: a meta-analysis. Ann Intern Med 1993;119:599-605.
- 2. Ackermann RT, Mulrow CD, Ramirez G et al. Garlic shows promise for improving some cardiovascular risk factors. Arch Intern Med 2001;161:813-824.
- Stevinson C, Pittler MH, Ernst E. Garlic for treating hypercholesterolemia: a meta-analysis of randomized clinical trials. Ann Intern Med 2000;133:420-429.
- 4. Silagy C, Neil A. Garlic as a lipid lowering agent—a meta-analysis. J R Coll Physicians Lond 1994;28:39-45.
- Lawson LD, Wang ZJ, Papadimitriou D. Allicin release under simulated gastrointestinal conditions from garlic powder tablets employed in clinical trials on serum cholesterol. Planta Med 2001;67:13-18.
- Lawson LD, Wang ZJ. Low allicin release from garlic supplements: a major problem due to the sensitivities of alliinase activity. J Agric Food Chem 2001;49:2592-2599.
- 7. Fleischauer AT, Poole C, Arab L. Garlic consumption and cancer prevention: meta-analyses of colorectal and stomach cancers. Am J Clin Nutr 2000;72:1047-1052.
- 8. El-Bayoumy K, Sinha R, Pinto JT, Rivlin RS. Cancer chemoprevention by garlic and garlic-containing sulfur and selenium compounds. J Nutr 2006;136:864S-869S.
- 9. Blumenthal M, Busse W, Goldberg A et al. The Complete German Commission E Monographs. Austin, Texas: American Botanical Council: Integrative Medicine Communications; 1998.
- 10. ESCOP. Allii Sativi Bulbus. ESCOP Monographs: the Scientific Foundation for Herbal Medicinal Products. 2nd ed. Exeter, UK: European Scientific Cooperative on Phytotherapy and Thieme; 2003:14-25.
- 11. WHO. Bulbus Allii Sativi. WHO Monographs on Selected Medicinal Plants. 1 vol. Geneva: World Health Organization; 1999:16-33.
- 12. McKenna D, Jones K, Hughes K, Humphrey S. Garlic. Botanical Medicines. 2nd ed. Binghampton: Haworth Press; 2002:375-409.
- 13. Vaes LP, Chyka PA. Interactions of warfarin with garlic, ginger, ginkgo, or ginseng: nature of the evidence. Ann Pharmacother 2000;34:1478-1482.
- 14. Steiner M, Li W. Aged garlic extract, a modulator of cardiovascular risk factors: a dose-finding study on the effects of AGE on platelet functions. J Nutr 2001;131:980S-984S.
- 15. Rahman K, Billington D. Dietary supplementation with aged garlic extract inhibits ADP-induced platelet aggregation in humans. J Nutr 2000;130:2662-2665.
- 16. Kiesewetter H, Jung F, Pindur G et al. Effect of garlic on thrombocyte aggregation, microcirculation, and other risk factors. Int J Clin Pharmacol Ther Toxicol 1991;29:151-155.
- 17. Kiesewetter H, Jung F, Jung EM et al. Effect of garlic on platelet aggregation in patients with increased risk of juvenile ischaemic attack. Eur J Clin Pharmacol 1993;45:333-336.
- 18. German K, Kumar U, Blackford HN. Garlic and the risk of TURP bleeding. Br J Urol 1995;76:518.
- 19. Rose KD, Croissant PD, Parliament CF, Levin MB. Spontaneous spinal epidural hematoma with associated platelet dysfunction from excessive garlic ingestion: a case report. Neurosurgery 1990;26:880-882.
- 20. Burnham BE. Garlic as a possible risk for postoperative bleeding. Plast Reconstr Surg 1995;95:213.
- 21. Macan H, Uykimpang R, Alconcel M et al. Aged garlic extract may be safe for patients on warfarin therapy. J Nutr 2006;136:793S-795S.
- 22. Siegel G, Walter A, Engel S et al. [Pleiotropic effects of garlic]. Wien Med Wochenschr 1999;149:217-224.
- 23. Thomson M, Al-Qattan KK, Bordia T, Ali M. Including garlic in the diet may help lower blood glucose, cholesterol, and triglycerides. J Nutr 2006;136:800S-802S.
- 24. Budoff M. Aged garlic extract retards progression of coronary artery calcification. J Nutr 2006;136:741S-744S.
- 25. Yeh YY, Yeh SM. Homocysteine-lowering action is another potential cardiovascular protective factor of aged garlic extract. J Nutr 2006;136:745S-749S.
- 26. Weiss N, Ide N, Abahji T et al. Aged garlic extract improves homocysteine-induced endothelial dysfunction in macro- and microcirculation. J Nutr 2006;136:750S-754S.
- 27. Borek C. Garlic reduces dementia and heart-disease risk. J Nutr 2006;136:810S-812S.
- 28. Lash JP, Cardoso LR, Mesler PM et al. The effect of garlic on hypercholesterolemia in renal transplant patients. Transplant Proc 1998;30:189-191.
- 29. Teyssier C, Guenot L, Suschetet M, Siess MH. Metabolism of diallyl disulfide by human liver microsomal cytochromes P-450 and flavin-containing monooxygenases. Drug Metab Dispos 1999;27:835-841.

- 30. Jin L, Baillie TA. Metabolism of the chemoprotective agent diallyl sulfide to glutathione conjugates in rats. Chem Res Toxicol 1997;10:318-327.
- 31. Wargovich MJ. Diallylsulfide and allylmethylsulfide are uniquely effective among organosulfur compounds in inhibiting CYP2E1 protein in animal models. J Nutr 2006;136:832S-834S.
- 32. Tanaka E, Terada M, Misawa S. Cytochrome P450 2E1: its clinical and toxicological role. J Clin Pharm Ther 2000;25:165-175.
- 33. Zhou S, Koh HL, Gao Y et al. Herbal bioactivation: the good, the bad and the ugly. Life Sci 2004;74:935-968.
- 34. Loizou GD, Cocker J. The effects of alcohol and diallyl sulphide on CYP2E1 activity in humans: a phenotyping study using chlorzoxazone. Hum Exp Toxicol 2001;20:321-327.
- 35. Gurley BJ, Gardner SF, Hubbard MA et al. Cytochrome P450 phenotypic ratios for predicting herb-drug interactions in humans. Clin Pharmacol Ther 2002;72:276-287.
- 36. Davenport D, Wargovich M. Modulation of cytochrome P450 enzymes by organosulfur compounds from garlic. Food Chem Toxicol 2005.
- 37. Yang CS, Chhabra SK, Hong JY, Smith TJ. Mechanisms of inhibition of chemical toxicity and carcinogenesis by diallyl sulfide (DAS) and related compounds from garlic. J Nutr 2001;131:1041S-1045S.
- 38. Taubert D, Glockner R, Muller D, Schomig E. The garlic ingredient diallyl sulfide inhibits cytochrome P450 2E1 dependent bioactivation of acrylamide to glycidamide. Toxicol Lett 2006;164:1-5.
- 39. Wang EJ, Li Y, Lin M et al. Protective effects of garlic and related organosulfur compounds on acetaminophen-induced hepatotoxicity in mice. Toxicol Appl Pharmacol 1996;136:146-154.
- 40. Milner JA. Preclinical perspectives on garlic and cancer. J Nutr 2006;136:827S-831S.
- 41. Herman-Antosiewicz A, Singh SV. Signal transduction pathways leading to cell cycle arrest and apoptosis induction in cancer cells by Allium vegetable-derived organosulfur compounds: a review. Mutat Res 2004;555:121-131.
- 42. Gurley BJ, Gardner SF, Hubbard MA et al. Clinical assessment of effects of botanical supplementation on cytochrome P450 phenotypes in the elderly: St John's wort, garlic oil, Panax ginseng and Ginkgo biloba. Drugs Aging 2005;22:525-539.
- 43. Marchand LL, Wilkinson GR, Wilkens LR. Genetic and dietary predictors of CYP2E1 activity: a phenotyping study in Hawaii Japanese using chlorzoxazone. Cancer Epidemiol Biomarkers Prev 1999;8:495-500.
- 44. Milner JA. Molecular targets for bioactive food components. J Nutr 2004;134:2492S-2498S.
- 45. Markowitz JS, Devane CL, Chavin KD et al. Effects of garlic (Allium sativum L.) supplementation on cytochrome P450 2D6 and 3A4 activity in healthy volunteers. Clin Pharmacol Ther 2003;74:170-177.
- 46. Foster BC, Foster MS, Vandenhoek S et al. An in vitro evaluation of human cytochrome P450 3A4 and P-glycoprotein inhibition by garlic. J Pharm Pharm Sci 2001;4:176-184.
- 47. Nabekura T, Kamiyama S, Kitagawa S. Effects of dietary chemopreventive phytochemicals on P-glycoprotein function. Biochem Biophys Res Commun 2005;327:866-870.
- 48. Greenblatt DJ, Leigh-Pemberton RA, von Moltke LL. In vitro interactions of water-soluble garlic components with human cytochromes p450. J Nutr 2006;136:806S-809S.
- 49. Matsuura H. Saponins in garlic as modifiers of the risk of cardiovascular disease. J Nutr 2001;131:1000S-1005S.
- 50. Manyike PT, Kharasch ED, Kalhorn TF, Slattery JT. Contribution of CYP2E1 and CYP3A to acetaminophen reactive metabolite formation. Clin Pharmacol Ther 2000;67:275-282.
- 51. Dooley TP. Molecular biology of the human phenol sulfotransferase gene family. J Exp Zool 1998;282:223-230.
- 52. Court MH, Duan SX, von Moltke LL et al. Interindividual variability in acetaminophen glucuronidation by human liver microsomes: identification of relevant acetaminophen UDP-glucuronosyltransferase isoforms. J Pharmacol Exp Ther 2001;299:998-1006.
- 53. Bernal W. Changing patterns of causation and the use of transplantation in the United Kingdom. Semin Liver Dis 2003;23:227-237.
- 54. Lee WM. Acute liver failure in the United States. Semin Liver Dis 2003;23:217-226.
- 55. Hu JJ, Yoo JS, Lin M et al. Protective effects of diallyl sulfide on acetaminophen-induced toxicities. Food Chem Toxicol 1996;34:963-969.
- 56. Hattori A, Yamada N, Nishikawa T et al. Protective effect of ajoene on acetaminophen-induced hepatic injury in mice. Biosci Biotechnol Biochem 2001;65:2555-2557.
- 57. Sumioka I, Matsura T, Yamada K. Therapeutic effect of S-allylmercaptocysteine on acetaminophen-induced liver injury in mice. Eur J Pharmacol 2001;433:177-185.
- 58. Sumioka I, Matsura T, Kasuga S et al. Mechanisms of protection by S-allylmercaptocysteine against acetaminophen-induced liver injury in mice. Jpn J Pharmacol 1998;78:199-207.
- 59. Lin MC, Wang EJ, Patten C et al. Protective effect of diallyl sulfone against acetaminophen-induced hepatotoxicity in mice. J Biochem Toxicol 1996;11:11-20.

- 60. Zhao C, Shichi H. Prevention of acetaminophen-induced cataract by a combination of diallyl disulfide and N-acetylcysteine. J Ocul Pharmacol Ther 1998;14:345-355.
- 61. Kalantari H, Salehi M. The protective effect of garlic oil on hepatotoxicity induced by acetaminophen in mice and comparison with N-acetylcysteine. Saudi Med J 2001;22:1080-1084.
- 62. Gwilt PR, Lear CL, Tempero MA et al. The effect of garlic extract on human metabolism of acetaminophen. Cancer Epidemiol Biomarkers Prev 1994;3:155-160.
- 63. Briggs WH, Xiao H, Parkin KL et al. Differential inhibition of human platelet aggregation by selected Allium thiosulfinates. J Agric Food Chem 2000;48:5731-5735.
- 64. Ali M, Bordia T, Mustafa T. Effect of raw versus boiled aqueous extract of garlic and onion on platelet aggregation. Prostaglandins Leukot Essent Fatty Acids 1999;60:43-47.
- 65. Apitz-Castro R, Badimon JJ, Badimon L. Effect of ajoene, the major antiplatelet compound from garlic, on platelet thrombus formation. Thromb Res 1992;68:145-155.
- 66. Apitz-Castro R, Badimon JJ, Badimon L. A garlic derivative, ajoene, inhibits platelet deposition on severely damaged vessel wall in an in vivo porcine experimental model. Thromb Res 1994;75:243-249.
- 67. Ariga T, Tsuji K, Seki T et al. Antithrombotic and antineoplastic effects of phyto-organosulfur compounds. BioFactors 2000;13:251-255.
- 68. Chan K-c, Hsu C-c, Yin M-c. Protective effect of three diallyl sulphides against glucose-induced erythrocyte and platelet oxidation, and ADP-induced platelet aggregation. Thromb Res 2002;108:317-322.
- 69. Qi R, Liao F, Inoue K et al. Inhibition by diallyl trisulfide, a garlic component, of intracellular Ca(2+) mobilization without affecting inositol-1,4,5-trisphosphate (IP[3]) formation in activated platelets. Biochem Pharmacol 2000;60:1475-1483.
- 70. Rendu F, Brohard-Bohn B, Pain S et al. Thiosulfinates inhibit platelet aggregation and microparticle shedding at a calpain-dependent step. Thromb Haemost 2001;86:1284-1291.
- 71. Allison GL, Lowe GM, Rahman K. Aged garlic extract and its constituents inhibit platelet aggregation through multiple mechanisms. J Nutr 2006;136:782S-788S.
- 72. Allison GL, Lowe GM, Rahman K. Aged garlic extract may inhibit aggregation in human platelets by suppressing calcium mobilization. J Nutr 2006;136:789S-792S.
- 73. Kiesewetter H, Jung F, Mrowietz C et al. Effects of garlic on blood fluidity and fibrinolytic activity: a randomised, placebo-controlled, double-blind study. Br J Clin Pract Suppl 1990;69:24-29.
- 74. Bordia A, Verma SK, Srivastava KC. Effect of garlic (Allium sativum) on blood lipids, blood sugar, fibrinogen and fibrinolytic activity in patients with coronary artery disease. Prostaglandins Leukot Essent Fatty Acids 1998;58:257-263.
- 75. Gadkari JV, Joshi VD. Effect of ingestion of raw garlic on serum cholesterol level, clotting time and fibrinolytic activity in normal subjects. J Postgrad Med 1991;37:128-131.
- 76. Apitz-Castro R, Escalante J, Vargas R, Jain MK. Ajoene, the antiplatelet principle of garlic, synergistically potentiates the antiaggregatory action of prostacyclin, forskolin, indomethacin and dypiridamole on human platelets. Thromb Res 1986;42:303-311.
- 77. Harenberg J, Giese C, Zimmermann R. Effect of dried garlic on blood coagulation, fibrinolysis, platelet aggregation and serum cholesterol levels in patients with hyperlipoproteinemia. Atherosclerosis 1988;74:247-249.
- 78. Teweleit S, Kuschel U, Hippius M et al. [Manifestation and prevention of adverse drug reactions (ADR) in the pharmacotherapy of cardiovascular diseases]. Med Klin 2001;96:442-450.
- 79. Pirmohamed M, James S, Meakin S et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18,820 patients. BMJ 2004;329:15-19.
- 80. Brouwer CA, Gietema JA, van den Berg MP et al. Long-term cardiac follow-up in survivors of a malignant bone tumour. Ann Oncol 2006;17:1586-1591.
- 81. Ewer M, Benjamin R. Cardiotoxcity of chemotherapeutic drugs. In: Perry M, ed. The Chemotherapy Source Book. 3rd ed. Philadelphia: Lippincott, Williams & Wilkins; 2001:458-468.
- 82. Kojima R, Toyama Y, Ohnishi ST. Protective effects of an aged garlic extract on doxorubicin-induced cardiotoxicity in the mouse. Nutr Cancer 1994;22:163-173.
- 83. Dwivedi C, John LM, Schmidt DS, Engineer FN. Effects of oil-soluble organosulfur compounds from garlic on doxorubicin-induced lipid peroxidation. Anticancer Drugs 1998;9:291-294.
- 84. Thabrew MI, Samarawickrema NA, Chandrasena LG, Jayasekera S. Protection by garlic against Adriamycin-induced alterations in the oxido-reductive status of mouse red blood cells. Phytother Res 2000;14:215-217.
- 85. Mukherjee S, Banerjee SK, Maulik M et al. Protection against acute Adriamycin-induced cardiotoxicity by garlic: role of endogenous antioxidants and inhibition of TNF-α expression. BMC Pharmacol 2003;3:16.
- 86. Borek C. Antioxidant health effects of aged garlic extract. J Nutr 2001;131:1010S-1015S.

- 87. Arora A, Seth K, Shukla Y. Reversal of P-glycoprotein-mediated multidrug resistance by diallyl sulfide in K562 leukemic cells and in mouse liver. Carcinogenesis 2004;25:941-949.
- 88. Chen C, Kong A-NT. Dietary chemopreventive compounds and ARE/EpRE signaling. Free Radical Biol Med 2004;36:1505-1516.
- 89. Gebhardt R. Multiple inhibitory effects of garlic extracts on cholesterol biosynthesis in hepatocytes. Lipids 1993;28:613-619.
- 90. Qureshi AA, Abuirmeileh N, Din ZZ et al. Inhibition of cholesterol and fatty acid biosynthesis in liver enzymes and chicken hepatocytes by polar fractions of garlic. Lipids 1983;18:343-348.
- 91. Gebhardt R. Inhibition of cholesterol biosynthesis by a water-soluble garlic extract in primary cultures of rat hepatocytes. Arzneimittelforschung 1991;41:800-804.
- 92. Liu L, Yeh YY. S-alk(en)yl cysteines of garlic inhibit cholesterol synthesis by deactivating HMG-CoA reductase in cultured rat hepatocytes. J Nutr 2002;132:1129-1134.
- 93. Kumar RV, Banerji A, Kurup CK, Ramasarma T. The nature of inhibition of 3-hydroxy-3-methylglutaryl CoA reductase by garlic-derived diallyl disulfide. Biochim Biophys Acta 1991;1078:219-225.
- 94. Singh DK, Porter TD. Inhibition of sterol  $4\alpha$ -methyl oxidase is the principal mechanism by which garlic decreases cholesterol synthesis. J Nutr 2006;136:759S-764S.
- 95. Silagy CA, Neil HA. A meta-analysis of the effect of garlic on blood pressure. J Hypertens 1994;12:463-468.
- 96. Langsjoen PH, Langsjoen AM. The clinical use of HMG CoA-reductase inhibitors and the associated depletion of coenzyme Q10: a review of animal and human publications. Biofactors 2003;18:101-111.
- 97. Lang A, Lahav M, Sakhnini E et al. Allicin inhibits spontaneous and TNF-α induced secretion of proinflammatory cytokines and chemokines from intestinal epithelial cells. Clin Nutr 2004;23:1199-1208.
- 98. Horie T, Matsumoto H, Kasagi M et al. Protective effect of aged garlic extract on the small intestinal damage of rats induced by methotrexate administration. Planta Med 1999;65:545-548.
- 99. Horie T, Awazu S, Itakura Y, Fuwa T. Alleviation by garlic of antitumor drug-induced damage to the intestine. J Nutr 2001;131:1071S-1074S.
- 100. Khosla P, Karan RS, Bhargava VK. Effect of garlic oil on ethanol induced gastric ulcers in rats. Phytother Res 2004;18:87-91.
- 101. Piscitelli SC, Burstein AH, Welden N et al. The effect of garlic supplements on the pharmacokinetics of saquinavir. Clin Infect Dis 2002;34:234-238.
- 102. Gallicano K, Foster B, Choudhri S. Effect of short-term administration of garlic supplements on single-dose ritonavir pharmacokinetics in healthy volunteers. Br J Clin Pharmacol 2003;55:199-202.
- 103. Patel J, Buddha B, Dey S et al. In vitro interaction of the HIV protease inhibitor ritonavir with herbal constituents: changes in P-gp and CYP3A4 activity. Am J Ther 2004;11:262-277.
- 104. Mouly SJ, Paine MF, Watkins PB. Contributions of CYP3A4, P-glycoprotein, and serum protein binding to the intestinal first-pass extraction of saquinavir. J Pharmacol Exp Ther 2004;308:941-948.
- 105. Gisolf EH, van Heeswijk RP, Hoetelmans RW, Danner SA. Decreased exposure to saquinavir in HIV-1-infected patients after long-term antiretroviral therapy including ritonavir and saquinavir. AIDS 2000;14:801-805.
- 106. Sunter WH. Warfarin and garlic [letter]. Pharmacol J 1991;246:722.
- 107. Fugh-Berman A, Ernst E. Herb-drug interactions: review and assessment of report reliability. Br J Clin Pharmacol 2001;52:587-595.
- 108. Miller LG. Herbal medicinals: selected clinical considerations focusing on known or potential drug-herb interactions. Arch Intern Med 1998;158:2200-2211.
- 109. Rozenfeld V, Sisca T, Callahan A, Crain J. Double-blind, randomized, placebo-controlled trial of aged garlic extract inpatients stabilized on warfarin therapy [poster]. American Society of Clinical Pharmacists, Mid-year Clinical Meeting. Las Vegas; 2000.
- 110. Pathak A, Leger P, Bagheri H et al. Garlic interaction with fluindione: a case report. Therapie 2003;58:380-381.
- 111. Comets E, Pousset F, Mentre F et al. Prediction of fluindione maintenance dosage hampered by large intraindividual variability. Ther Drug Monit 2000;22:668-675.
- 112. Loebstein R, Yonath H, Peleg D et al. Interindividual variability in sensitivity to warfarin—nature or nurture? Clin Pharmacol Ther 2001;70:159-164.
- 113. Fitzmaurice DA, Blann AD, Lip GY. Bleeding risks of antithrombotic therapy. BMJ 2002;325:828-831.
- 114. Wells PS, Holbrook AM, Crowther NR, Hirsh J. Interactions of warfarin with drugs and food. Ann Intern Med 1994;121:676-683.
- 115. Borgert CJ, Borgert SA, Findley KC. Synergism, antagonism, or additivity of dietary supplements: application of theory to case studies. Thromb Res 2005;117:123-132; discussion 145-151.
- 116. Marder VJ. The interaction of dietary supplements with antithrombotic agents: scope of the problem. Thromb Res 2005;117:7-13; discussion 39-42.

- 117. Ciplea AG, Richter KD. The protective effect of Allium sativum and crataegus on isoprenaline-induced tissue necroses in rats. Arzneimittelforschung 1988;38:1583-1592.
- 118. Saxena KK, Gupta B, Kulshrestha VK et al. Effect of garlic pretreatment on isoprenaline-induced myocardial necrosis in albino rats. Indian J Physiol Pharmacol 1980;24:233-236.
- 119. Banerjee SK, Sood S, Dinda AK et al. Chronic oral administration of raw garlic protects against isoproterenol-induced myocardial necrosis in rat. Comp Biochem Physiol C Toxicol Pharmacol 2003;136:377-386.
- 120. Brinker F. Herb Contraindications and Drug Interactions. 3rd ed. Sandy, Ore: Eclectic Medical Publications; 2001.
- 121. Martin N, Bardisa L, Pantoja C et al. Experimental cardiovascular depressant effects of garlic (Allium sativum) dialysate. J Ethnopharmacol 1992;37:145-149.
- 122. Martin N, Bardisa L, Pantoja C et al. Anti-arrhythmic profile of a garlic dialysate assayed in dogs and isolated atrial preparations. J Ethnopharmacol 1994;43:1-8.
- 123. Martin N, Bardisa L, Pantoja C et al. Involvement of calcium in the cardiac depressant actions of a garlic dialysate. J Ethnopharmacol 1997;55:113-118.
- 124. Jain RC, Vyas CR. Garlic in alloxan-induced diabetic rabbits. Am J Clin Nutr 1975;28:684-685.
- 125. Mathew PT, Augusti KT. Studies on the effect of allicin (diallyl disulphide-oxide) on alloxan diabetes. I. Hypoglycaemic action and enhancement of serum insulin effect and glycogen synthesis. Indian J Biochem Biophys 1973;10:209-212.
- 126. Zhang XH, Lowe D, Giles P et al. Gender may affect the action of garlic oil on plasma cholesterol and glucose levels of normal subjects. J Nutr 2001;131:1471-1478.
- 127. Huang CN, Horng JS, Yin MC. Antioxidative and antiglycative effects of six organosulfur compounds in low-density lipoprotein and plasma. J Agric Food Chem 2004;52:3674-3678.
- 128. Ahmad MS, Ahmed N. Antiglycation properties of aged garlic extract: possible role in prevention of diabetic complications. J Nutr 2006;136:796S-799S.